



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C07F 15/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 91/03482 (43) International Publication Date: 21 March 1991 (21.03.91)</p>
<p>(21) International Application Number: PCT/US90/04966 (22) International Filing Date: 4 September 1990 (04.09.90) (30) Priority data: 401,919 1 September 1989 (01.09.89) US (71) Applicant: THE UNIVERSITY OF VERMONT AND STATE AGRICULTURAL COLLEGE [US/US]; Burlington, VT 05405-3160 (US). (72) Inventor: FARRELL, Nicholas ; 306 Woolen Mill, Winoski, VT 05405 (US). (74) Agent: HUNTINGTON, R., Danny; Burns, Doane, Swecker & Mathis, The George Mason Building, Washington & Prince Streets, P.O. Box 1404, Alexandria, VA 22313-1404 (US).</p>		<p>(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent)*, DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>

(54) Title: NEW BIS-PLATINUM COMPLEXES AS CHEMOTHERAPEUTIC AGENTS



(57) Abstract

A bis(platinum) complex having structure (I), wherein X, Y, Z, X', Y', and Z' are the same or different ligands and are halide, sulphate, phosphate, nitrate, carboxylate, substituted carboxylate, dicarboxylate, substituted dicarboxylate primary or secondary amine, sulfoxide, phosphine, pyridine-type nitrogen and the like; preferably, there is at least one Pt-anion bond on each Pt molecule; and A is a diamine or polyamine. Complexes of trans geometry [PtX₂(L)(L')] where at least L is a planar ligand such as pyridine but preferably quinoline and L' may be amine, sulfoxide or substituted sulfoxide and X is a chloride, nitrate or carboxylate. A process for obtaining bis(platinum) complexes where the two platinum coordination spheres may contain different ligands in each sphere and where the geometry of each platinum coordination sphere may be the same or different.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Mosaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
DE	Germany	LU	Luxembourg	TD	Chad
DK	Denmark			TG	Togo
				US	United States of America

WO 91/03482

PCT/US90/04966

-1-

NEW BIS-PLATINUM COMPLEXES
AS CHEMOTHERAPEUTIC AGENTS

BACKGROUND OF THE INVENTION

5 The present invention relates to novel
bis(platinum) complexes and to pharmaceutical
compositions containing them.

 The use of platinum complexes in cancer
chemotherapy is well known. A number of platinum
10 complexes, such as Platinol, a registered trademark of
cisplatin manufactured by Bristol Myers, Co., are used
to treat testicular, ovarian, head and neck, and small-
cell lung carcinomas. However, treatment with
cisplatin may result in severe nephrotoxicity. A
15 further clinical disadvantage is the problem of
acquired drug resistance resulting in the tumor
becoming refractory to treatment by the agent.

 To overcome the nephrotoxic effects of
cisplatin, a second-generation analog, carboplatin, was
20 developed. Paraplatin is a registered trademark for
carboplatin manufactured by Bristol-Myers, Co.
Carboplatin, or $[\text{Pt}(\text{NH}_3)_2(\text{CBDCA})]$ (where CBDCA is
1,1'-cyclobutanedicarboxylate), is effective against the
same spectrum of carcinomas as cisplatin, but exhibits
25 a marked reduction in the nephrotoxic effects.

 A number of different platinum compounds have
been developed in an attempt to treat different tumors
or carcinomas. For instance, U.S. Patent No. 4,225,529
discloses the use of a cis coordination compound of
30 platinum having four ligands which are selected from
the group consisting of halides, sulphates,
phosphates, nitrates, carboxylates, and same or
different straight-chain amines which are coordinated
to the platinum atom through their nitrogen atoms.
35 These complexes are utilized for treating L-1210
leukemia in mice.

WO 91/03482

PCT/US90/04966

-2-

Also, U.S. Patent Nos. 4,250,189, 4,553,502, and 4,565,884 relate to various Pt(II) and Pt(IV) complexes having antitumor activity. These bis(platinum) complexes are linked with a carboxylate linkage such that upon administration of these complexes to the patient, the complexes undergo rapid hydrolysis to produce two cis monoplutonium moieties which are then delivered to the active site.

However, in U.S. Patent No. 4,797,393, a bis(platinum) complex is disclosed, which complex is delivered intact to the active site. This bis(platinum) complex has a bridging diamine or polyamine ligand and has primary or secondary amines or pyridine type nitrogens attached to the platinum complex, as well as two different or identical ligands which may be a halide, sulphate, phosphate, nitrate, carboxylate, substituted carboxylate or dicarboxylate.

Most of the synthesis of platinum analogs to date has been based on the [cis-Pt(amine)₂X₂] structure where X is a chloride or an anionic leaving group since it is believed that the cis configuration is necessary for antitumor activity in monomeric platinum complexes. A wide range of amines has been employed and a major emphasis has been on 1,2-diaminocyclohexane (which is oftentimes referred to as "dach") because laboratory studies (Burchenal et al. *Biochimie*, 1978, 60, 961) show that complexes derived from these amines are non-cross-resistant with cisplatin. This means that dach complexes maintain their curative activity in tumor cell lines resistant to cisplatin and the clinical advantage of such an agent should be apparent. The mechanism of action of cisplatin is generally believed to be by formation of crosslinks, especially interstrand crosslinks on DNA, producing an overall conformational change on the DNA, which eventually leads to the inhibition of replication

WO 91/03482

PCT/US90/04966

-3-

and thus produces a cytotoxic effect as discussed in Sherman and Lippard, Chem. Review, 1987, 87, 1153 and Reedijk et al, Structure and Bonding, 1987, 67, 53.

Even though other closely related platinum complexes such as those in the trans-configuration [trans-Pt(NH₃)₂X₂], trans-DDP, and monodentate complexes [(Pt(NH₃)₃Cl]⁺ and [Pt(dien)Cl]⁺, (dien = diethylenetriamine, a tridentate amine) do bind to DNA, they do not exhibit antitumor activity. This is because the trans-[Pt(NH₃)₂Cl₂] and especially monodentate species such as [Pt(NH₃)₃Cl]⁺ cannot form the 1,2-intrastrand crosslink.

It has been discovered that bis(platinum) complexes of U.S. Patent 4,797,393 may exhibit high cytotoxic activity and are non-cross-resistant with both cisplatin and platinum-dach complexes. Their DNA binding involves interstrand crosslinks formed because of the bimetallic nature of the bis(platinum) complexes whereby each platinum atom of the bis(platinum) unit binds to opposite DNA strands. This effect has led in part to the discovery of structurally new bis(platinum) derivatives which also have activity in cisplatin-resistant lines and thus may have a broader spectrum of activity than cisplatin. Thus, there remains a need in the art to produce pharmaceutical compounds which are active in cisplatin-resistant lines.

SUMMARY OF THE INVENTION

Accordingly, it is an object of the present invention to provide a process for producing stable bis(platinum) complexes to be used for antitumor and pharmaceutical applications.

It is another object of the present invention to produce bis(platinum) complexes that induce interstrand cross-links to the DNA molecule.

WO 91/03482

PCT/US90/04966

-4-


It is yet another object of the present invention to provide a bis(platinum) complex having at least one Pt-Cl bond on each platinum, such that each Pt atom of the bis(platinum) molecule may bind to opposite strands of DNA.

A still further object of the present invention is to provide a bis(platinum) complex in cis or trans isomeric form that exhibits antitumor activity.

A further object is to demonstrate the activation of monomeric trans complexes which exhibit cytotoxicity equivalent to cisplatin and which may exhibit enhanced antitumor activity by themselves or upon incorporation into the new bis(platinum) structures.

A further object is to demonstrate a method for obtaining bis(platinum) complexes where the two platinum coordination spheres are different by using as a precursor a monomeric platinum complex containing only one end of a diamine bound to the platinum.

In accordance with the foregoing objectives, the present invention provides a bis(platinum) complex having the general formula:



wherein X, Y, Z, X', Y' and Z' may be a combination of anionic groups or neutral groups and A is a bridging ligand.

These and other objects, features and advantages will be apparent from the following more particular description of the preferred embodiments.

35

WO 91/03482

PCT/US90/04966

-5-


BRIEF DESCRIPTION OF THE DRAWING

The Figure depicts the formation of an interstrand crosslink through binding of one Pt of the bis(platinum) molecule to at least one base on one strand of DNA with concomitant binding of the other Pt atom to a base on the other strand.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention relates to novel bis(platinum) complexes and to the preparation of these complexes. These novel platinum complexes have coordination spheres which may be monodentate and thus contain only one anionic group, such as a chloride ion, attached to each platinum atom or the platinum coordination spheres may be different and may contain, for example, either one or two anionic groups, such as chloride, in each sphere. Further, the geometry of the coordination spheres may be cis or trans or a mixture thereof; that is, one coordination sphere may be cis and the other may be trans.

A general formula of the bis(platinum) complexes encompassed in the present invention include:



wherein X, Y, Z, X', Y', and Z' may be a combination of anionic groups such as halides including chlorine, bromine, iodine and fluorine and pseudohalides, as well as sulphate, phosphate, phosphonate, nitrate, carboxylate, substituted carboxylated, dicarboxylate and substituted dicarboxylate, or any neutral group such as a primary or secondary amine, sulfoxide (such as DMSO), phosphine, pyridine, or planar aromatic or

WO 91/03482

PCT/US90/04966

-6-


pseudo-aromatic pyridine-like ligand such as substituted pyridine, quinoline, imidazole, thiazole, pyrimidine, purine, acridine, pyrazole, benzimidazole, benzothiazole and the like, as well as sulfoxide and phosphine. It is preferable to use at least one chloride group per platinum atom. Where there is only one anionic group attached to each Pt atom, each Pt atom is monodentate and the complexes carry an overall 2⁺ charge. Where there is one anionic group attached to one Pt atom and two to the second Pt atom; the complexes therefore carry a 1⁺ charge, with one Pt atom monodentate and the other bidentate.

The bridging ligand A is a diamine or polyamine wherein the primary amine N atoms are coordinated to the Pt atom such that platinum is present as Pt²⁺ and preferably has the formula:

$$\text{NDE}-(\text{C}(\text{R}^4)_2)_r-(\text{R}^5)_o-(\text{C}(\text{R}^6)_2)_p-\text{NDE}$$

in which r and p are integers from 1 to 4, inclusive and o is 0 or 1; and the R⁴ and R⁶ groups are the same or different and are hydrogen, lower alkyl, aryl, cycloalkyl, cycloalkenyl, aralkyl, halogen, pseudohalogen, hydroxy, alkoxy, aryloxy, carboxylic acid ester, or carboxylic acid salt. Preferably, all R⁴ and R⁶ groups are hydrogen.

The R⁵ group is optional, and if employed is selected from alkyl, aryl (such as phenyl), amino, alkylamino, diamino of the formula:



wherein q is an integer of 1 to 4, inclusive, hydroxyalkyl, alkoxy, sulfur or oxygen.

The D and E groups are the same or different and are selected from hydrogen, lower alkyl, aryl, alkaryl, aralkyl, alkenyl, cycloalkyl, cycloalkenyl, halogen, pseudohalogen, hydroxy, alkoxy, aryloxy, or


WO 91/03482

PCT/US90/04966

-7-


sulphonic acids or salts thereof. The preferred substituent is hydrogen.

Particularly preferred A bridging ligands include straight chain diamines. Such bis(platinum) complexes preferably have the formula:



wherein X, Y, Z, X', Y' and Z' may each be represented by the groups set forth above and n is represented by 2 to 20 and preferably 4 to 12, inclusive.

The sulfoxide preferably has the structure



The amines may be aliphatic or aromatic and generally include ammonia, branched or straight chain lower alkyl amines, aryl amines, aralkyl amines, lower alkenyl amines, cycloalkyl amines, cycloalkenyl amine, and polycyclic hydrocarbon amines.

Substituted or unsubstituted heterocyclic amines, nucleosides, nucleotides, pyridine-type nitrogen containing compounds, and the like may be used in the practice of the present invention. Suitable substituents include but are not limited to alkyl, aromatic aryl, hydroxy, lower alkoxy, carboxylic acid or acid ester, nitro and halogen substituents.

Purines and pyrimidines which are suitable in the practice of the present invention include, for example, cytosine, uracil, thymine, guanine, adenine,


WO 91/03482

PCT/US90/04966

-8-

xanthine, hypoxanthine, purine, pyrimidine and their substituted derivatives.

Where the anionic group is a carboxylate or a substituted carboxylate, the anionic group may be represented by the formula:



wherein m is an integer from 0 to 5, inclusive. The R³ groups may be the same or different and may be hydrogen, substituted or unsubstituted straight or branched chain alkyl, aryl, alkaryl, alkenyl, cycloalkyl, cycloalkenyl, halogen, pseudohalogen, hydroxy, carbonyl, formyl, nitro, amido, amino, alkoxy, aryloxy, sulphonic acid salt, carboxylic acid ester or carboxylic acid salt. Furthermore, the R³ groups can be combined so that two R³ groups represent a double bond oxygen or sulphur atom.

Lower alkyl and lower alkenyl in the present specification means one to five carbon atoms. Unless indicated otherwise, alkyl or alkenyl means 1 to 12 carbon atoms. By cycloalkyl is meant chains of 3 to 10 carbon atoms. Substituted in the present specification, unless indicated otherwise, is intended to mean substitution with a group chosen from alkyl, aryl, cycloalkyl of 3 to 10 carbon atoms, cycloalkenyl, aralkyl, halogen, pseudohalogen, hydroxy, alkoxy, cycloamino, or carboxylic acid salts or esters of one to ten carbon atoms.

The term pseudohalide in the present invention has the meaning found on page 560 of "Advanced Inorganic Chemistry" by Cotton and Wilkinson, Interscience Publishers, 1966. The text describes a pseudohalogen, i.e., pseudohalide, as being a molecule consisting of more than two electronegative atoms which, in the free state, represent halogen atoms.

WO 91/03482

PCT/US90/04966

-9-

Examples of these molecules include cyanide, cyanate, thiocyanate, and azide.

Preferably, there are one or two chlorine atoms on each Pt atom; thus, a total of two to four chlorine atoms are present on the preferred compounds of the present invention.

The bis(platinum) complexes of the present invention may contain a platinum moiety in the trans configuration, as well as the monodentate bis(platinum) complex. These complexes differ from the monomeric platinum complexes in that the mechanism of binding to DNA is different. The principal adducts of cisplatin, for example, are the intrastrand crosslinks between adjacent guanines (GG), and between a guanine-adenine neighboring pair (GA). The cytotoxic effect is believed to be induced by the conformational change of the DNA due to the Pt binding. Complexes such as trans-[Pt(NH₃)₂Cl₂] and the monodentate species such as [Pt(NH₃)₃Cl]⁺ have no cytotoxic effect presumably because their binding precludes any intrastrand crosslinking, even though they bind avidly to the DNA molecule.

In contrast to monomeric complexes, the bis(platinum) derivatives of U.S. Patent 4,797,393 are able to bind to the DNA molecule by two principal methods: an intrastrand crosslink formed in the same manner as described above and also an interstrand crosslink caused by each unique Pt atom binding to the two strands of the DNA molecule. See the Figure. The schematic depicts four base pairs, however, the base pair separation is schematic and is not intended to be restricted to the separation depicted. In the Figure, L and L' are any ligands on the platinum which are not directly involved with the initial interstrand crosslink formation. The complexes of the present invention cannot, by nature of their structure, form

WO 91/03482

PCT/US90/04966

-10-

intrastrand links on both Pt atoms but the agents remain highly cytotoxic, especially in cisplatin-resistant lines. This is despite the fact that the monomeric precursors of monodentate or trans geometry exert no cytotoxic effect. The combination of these units, however, into a bis(platinum) complex of the general formula disclosed herein, results in significant cytotoxicity. The dominant and uniting feature of all these new bis(platinum) complexes is their ability to induce interstrand crosslinking and the formation of at least one Pt-nucleic acid base bond on each strand of DNA. These features appear to be a sufficient requisite for cytotoxicity. Thus, the bimetallic agents forming interstrand crosslinking between the DNA strands are excellent chemotherapeutic agents, especially for those tumors which are resistant to cisplatin.

The bis(platinum) complexes according to the present invention are intended for pharmaceutical application. The complex is useful for the identical diseases and modalities and use in the same patients as cisplatin. This includes the treatment of tumors, radiation sensitization or potentiation (Douple et al, Cisplatin Current Status and New Developments, Eds. A. W. Prestayko, S.T. Crooke, and S.K. Carter, Academic Press, 125 (1980); Douple, Platinum Metals Rev., 1985, 29, 118), and parasitic diseases such as sleeping sickness (Farrell et al, Biochem. Pharmacology, 1984, 33, 961). The complexes of the present invention are administered at approximately the same dosage levels as cisplatin, while taking into account their LD₅₀ values. The complex is normally associated with a suitable pharmaceutical carrier. For example, the complex and carrier can be formulated for parenteral or oral administration by methods known in the art. For instance, see Remington's Pharmaceutical Sciences for


WO 91/03482

PCT/US90/04966


-11-

suitable pharmaceutically acceptable carriers and formulation methods.

The present invention also discloses a process for producing a bis(platinum) complex by use of monomeric precursors which contain a diamine or polyamine bound to the platinum atom by one of the amines, the other being uncomplexed (free or dangling). Reaction of these precursors with additional molecules containing at least one chloride ion capable of being displaced will give bis(platinum) complexes whose structure will depend on both the exact structure of the precursor molecule and the molecule to be appended. Specific examples are those of Examples 3 and 4 set forth below. The bridging and therefore bis(platinum) complex formation occurs through substitution by the free amino end of the precursor molecule of a Pt-Cl bond on the appended molecule. The precursor molecules are most simply represented as RNH_3^+Cl and reaction with the selected molecule gives the reaction:



where RNH_3 represents $PtX'Y'Z'A$ wherein Z' , Y' , Z' and A as well as X , Y and Z are as defined above. In the case of Example 3, the reaction is:



where R is $trans-[PtCl_2(NH_3)_2N(CH_2)_4]-$. See Reaction Scheme 2.

WO 91/03482

PCT/US90/04966

-12-

The reaction is preferably carried out in aqueous or methanolic solution in presence of a base. In the case of Example 3, the product precipitates from solution, is filtered off and recrystallized by standard methods known in the art.

In order to fully illustrate the present invention and the advantages thereof, the following specific examples are given, it being understood that these examples are intended only to be illustrative without serving as a limitation on the scope of the present invention.

The examples describe the preparation of the new bis(platinum) complexes and precursors for new bis(platinum) complexes by reaction with the starting materials of $\text{cis-[PtCl}_2(\text{NH}_3)_2]$ and $\text{K[PtCl}_3(\text{NH}_3)]$ respectively. The $\text{cis-[PtCl}_2(\text{NH}_3)_2]$ compound is prepared by the general method of Dhara, J. Indian Chem., 1970, 8, 913, while the $\text{K[PtCl}_3(\text{NH}_3)]$ compound is prepared by the general method of Abrams et al, Inorg. Chim. Acta, "A Convenient preparation of the Amminetri-chloroplatinate(II) Anion" 1987, 3, 131.

EXAMPLE 1

Preparation of $\{[\text{trans-[PtCl(NH}_3)_2)_2\text{H}_2\text{N(CH}_2)_n\text{NH}_2]\text{Cl}_2$

Monomeric $\text{trans-[PtCl}_2(\text{NH}_3)_2]$ was prepared from the tetra-amine complex by reacting $\text{cis-[PtCl}_2(\text{NH}_3)_2]$ with aqueous ammonia (NH_3) to form an intermediate, $[\text{Pt(NH}_3)_4]\text{Cl}_2$. This intermediate was further reacted with an aqueous hydrochloric acid solution to form $\text{trans-[PtCl}_2(\text{NH}_3)_2]$. $\text{Trans-[PtCl}_2(\text{NH}_3)_2]$ was reacted with 1,4-butanediamine in H_2O and stirred overnight. The clear solution was then filtered, evaporated to dryness and recrystallized from $\text{H}_2\text{O/acetone}$ or $\text{H}_2\text{O/EtOH}$ to give $\{[\text{trans-[PtCl(NH}_3)_2)_2\text{H}_2\text{N(CH}_2)_4\text{NH}_2]\text{Cl}_2$. The complex was characterized by elemental analysis as noted in Table 1

WO 91/03482

PCT/US90/04966

-13-

below, ^1H NMR (rel. to TMS) at 2.74 and 1.79 ppm and ^{195}Pt NMR at -2436.7 ppm rel. to PtCl_6^{2-} . The IR spectrum (KBr disc) shows bands typical of bridging diamine and $\nu(\text{Pt}-\text{Cl}) = 330\text{ cm}^{-1}$.

5 Examples of other trans complexes which may be used in this way include species such $[\text{PtCl}_2(\text{pyridine})_2]$, $\text{trans}-[\text{PtCl}_2(\text{pyridine})(\text{NH}_3)]$, $\text{trans}-[\text{PtCl}_2(\text{pyridine})(\text{DMSO})]$, $[\text{PtCl}_2(\text{quinoline})(\text{DMSO})]$, $[\text{PtCl}_2(\text{iso-quinoline})(\text{DMSO})]$.
10 The present inventor has also disclosed that the presence of a planar ligand such as pyridine or quinoline dramatically improves the cytotoxicity of complexes of the trans configuration. Table III presents the in vitro data with comparison of some NH_3
15 complexes. As will be seen, a complex such as $\text{trans}-[\text{PtCl}_2(\text{pyridine})_2]$ is in fact as cytotoxic as cisplatin and represents a further class of complexes non-cross-resistant with the parent cisplatin.

20 Thus, a further extension of the present discovery is the demonstration of the activation of trans complexes by use of planar ligands rather than NH_3 . The general formula for cytotoxic trans complexes with planar ligands is $[\text{PtX}_2(\text{L})(\text{L}')]]$ where X may be any anionic group and L and L' are the planar ligand such
25 as pyridine, quinoline, isoquinoline, acridine, pyrazole, thiazole, imidazole, benzimidazole, benzothiazole, and other pyridine-like planar aromatic or pseudo-aromatic heterocycles. Where L is not the same as L' and L represents the planar ligand as above,
30 then L' may represent a primary or secondary amine such as NH_3 or a sulfoxide such as DMSO. Incorporation of these structures into bis(platinum) complexes will occur in the same manner as outlined for $\text{trans}-[\text{PtCl}_2(\text{NH}_3)_2]$.
35

WO 91/03482

PCT/US90/04966

-14-

EXAMPLE 2Preparation of Trans-[PtCl₂(NH₃)(H₂N(CH₂)₄NH₃)]Cl

This example demonstrates the preparation of a metal complex containing only one end of the diamine initially bound. The preparative scheme for the precursor is outlined in Reaction Scheme 1 and is adapted from Farrell and Qu, Inorg. Chem., Chemistry of Bis(platinum) Complexes, "Formation of trans derivatives from tetra-amine complexes," In press, September 1989. The precursor, trans-[PtCl₂(NH₃)H₂N(CH₂)₄NH₃)]Cl was prepared by Steps 1 and 2 of Scheme 1 and the precursor complex is Product 2c. Product 2c contains one end of the diamine bound to the platinum and the other end free (or dangling). The free or dangling end of the diamine may then be used to produce new bis(platinum) complexes, as shown in Reaction Scheme 2. In Reaction Scheme 2, the free amine end is used to bind to another platinum atom but it may also be used to bind another metal.

The preparation is outlined below:

Step 1 of Reaction Scheme 1. 0.6 grams or 2 mmol of cis-[PtCl₂(NH₃)₂] was suspended in 20 ml of water and 0.177 grams or 2 mmol of 1,4-diaminobutane was added. This mixture was then stirred at 60°C for 1-1.5 hours. The solution was then filtered and evaporated to 1 ml. The product was precipitated by refrigeration at 3°C for about 24 hours. The precipitated product was then filtered, washed with EtOH and dried. The complex was further recrystallized from H₂O/EtOH. The yield was about 68% for the product [(cis-Pt(NH₃)₂(H₂N(CH₂)₄NH₂))₂]Cl₄ (Compound 2a).

Step 2 of Reaction Scheme 1) 0.5 grams of Complex 2a, [(cis-Pt(NH₃)₂(H₂N(CH₂)₄NH₂))₂]Cl₄ or 0.64 mmol was dissolved in 2 ml H₂O and 50 ml 6N HCl was then added. The solution was allowed to react for 6-8 hours at 60-70°C and was stirred constantly during that

WO 91/03482

PCT/US90/04966

-15-


period. A yellow solid precipitate then formed, which was filtered off and washed with H₂O/acetone and further dried in vacuo. The filtrate was recrystallized from DMA/0.1 N HCl. The yield for the product [(trans-PtCl₂(NH₃))₂H₂N(CH₂)₄NH₂] was about 48% (Product 2b). The IR spectrum showed bands at $\nu(\text{NH}) = 3280, 3235, 3195 \text{ cm}^{-1}$, $\nu(\text{Pt-Cl}) = 340 \text{ cm}^{-1}$. The ¹H NMR in d₇-DMF gave peaks at 1.63, 2.68 ppm and the ¹⁹⁵Pt NMR gave a peak at -2167 ppm.

WO 91/03482

PCT/US90/04966

-16-

REACTION SCHEME 1




WO 91/03482

PCT/US90/04966

-17-

REACTION SCHEME 2



WO 91/03482

PCT/US90/04966

-18-

The trans-[PtCl₂(NH₃)(H₂N(CH₂)₄NH₃)]Cl complex was isolated by evaporation of the above filtrate to an oil, upon filtering off Product 2b. Approximately 30 ml of EtOH was then added to the oil and the solution was stirred for approximately 30 minutes. A yellow solid was formed and filtered off. The filtrate was then washed with EtOH and dried. Trans-[PtCl₂(NH₃)(H₂N(CH₂)₄NH₃)]Cl was then recrystallized from H₂O/EtOH (Product 2c). The IR spectrum shows bands corresponding to $\nu(\text{NH})$ at 3290(sh), 3245, 3200 and $\nu(\text{Pt-Cl})$ at 335 cm⁻¹. The ¹H NMR in D₂O gave peaks at 1.8, 2.72(t) and 3.1 ppm. The ¹³C NMR in D₂O gave four peaks at 47.8, 42.0, 29.6 and 26.8 ppm. The ¹⁹⁵Pt NMR gave a peak at -2132 ppm.

EXAMPLE 3

Formation of bis(platinum) complexes with the platinum coordination spheres in different geometries.

Preparation of [(trans-PtCl₂(NH₃))H₂N(CH₂)_nNH₂(cis-PtCl₂(NH₃))]

This procedure demonstrates the use of a metal complex containing only one end of the diamine initially bound and the subsequent platination of the free or dangling amine end to produce bis(platinum) complexes (Reaction Scheme 2). To the precursor, trans-[PtCl₂(NH₃)(H₂N(CH₂)₄NH₃)]Cl (Compound 2c), one equivalent of K[PtCl₃(NH₃)] in MeOH was added in the presence of triethylamine. The complex of [(cis-PtCl₂(NH₃))H₂N(CH₂)₄NH₂(trans-PtCl₂(NH₃)))] precipitated out of solution, was filtered, washed with water and acetone and dried. The new complex was characterized by elemental analysis: ¹H NMR and ¹⁹⁵Pt NMR (-2165 ppm and -2171 ppm).

WO 91/03482

PCT/US90/04966

-19-

EXAMPLE 4

Formation of bis(platinum) complexes
with two different coordination spheres

Preparation of [trans-(PtCl₂(Me₂SO)(H₂N(CH₂)₄NH₂)trans-
[PtCl₂(NH₃)]

Following the same procedure set forth in
Example 2, the precursor, trans-
[PtCl₂(NH₃)(H₂N(CH₂)₄NH₃)]Cl, was prepared as a source
of an amine for binding. This precursor was then
reacted with one equivalent of K[PtCl₃(Me₂SO)] anion in
MeOH in the presence of triethylamine. The precipitate
was then filtered, washed with water and acetone, and
dried. The specific structure formed from this
procedure was [trans(PtCl₂(Me₂SO)(H₂N(CH₂)₄NH₂)trans-
[PtCl₂(NH₃)]]. The new complex was characterized by
elemental analysis, IR spectrum $\nu(\text{Pt-Cl}) = 330$, $\nu(\text{SO}) =$
1115, $\nu(\text{NH}) = 3260, 3200, 3110 \text{ cm}^{-1}$, $^1\text{H NMR} = 3.5,$
1.85, and 2.65 ppm $^{195}\text{Pt NMR}(-2172 \text{ and } -3131 \text{ ppm})$.

Summary of Properties of NewBis(platinum) Complexes

The complexes as prepared in the preceding
examples were readily soluble in dimethylsulfoxide
(DMSO), dimethylformamide (DMF), and had good water
solubility when isolated as cationic complexes. The
data in Table I below summarizes the elemental analyses
obtained for the complexes derived from the procedures
set forth above.

WO 91/03482

PCT/US90/04966

-20-

TABLE I

ELEMENTAL ANALYSES
FOUND (CALCULATED)

	EXAMPLE	%C	%H	%N	%Cl
5	1	7.76(7.74)	3.61(3.68)	12.0(12.04)	20.11(20.32)
10	3	7.13(7.34)	2.63(2.77)	8.49(8.56)	21.79(21.68)
	4	10.59(10.07)	2.93(2.96)	5.65(5.87)	20.40(19.83)

15

EXAMPLE 5

Biological Activity

The monodentate bis(platinum) complexes, Example 1, and the platinum moiety containing one trans configuration on one Pt atom, and a cis anion configuration on the other Pt atom, Example 3, were tested for cytotoxic activity in various L-1210 murine leukemia cell lines. The tests were carried out in vitro according to the procedures outlined by M. P. Hacker et al in Cancer Research, 1985, 45, 4748. The term ID₅₀ refers to the concentration required to inhibit cell growth by 50%, and thus the lower the number, the more effective cytotoxin that particular agent. The bis(platinum) complexes of the present invention were compared to bis(platinum) complexes of equal chain length having only cis configurations and a typical monodentate monomeric complex [Pt(dien)Cl]Cl. Table II summarizes the results obtained.

WO 91/03482

PCT/US90/04966

-21-

TABLE II

<u>CYTOTOXICITY DATA</u>					
5	COMPLEX	L-1210/0	L-1210/DDP	L-1210/dach	Solvent
		ID ₅₀ uM	ID ₅₀ uM	ID ₅₀	
	Example 1	3.4	0.9	5.4	H ₂ O
	Example 3	0.76	3.36	2.6	10% DMSO
10	BisPt4	0.28	2.19	0.336	DMF
	BisPtmal	0.93	5.03	2.55	H ₂ O
	[Pt(dien)Cl]Cl	>20	>20	>20	H ₂ O
15	wherein Example 1 = [(trans-				
				[PtCl(NH ₃)] ₂ H ₂ N(CH ₂) ₄ NH ₂]Cl ₂	
	Example 3	=	[(cis-		
			PtCl ₂ (NH ₃)] ₂ H ₂ N(CH ₂) ₄ NH ₂ (trans-		
			PtCl ₂ (NH ₃))]		
	BisPt4	=	[(cis-PtCl ₂ (NH ₃)] ₂ (H ₂ N(CH ₂) ₄ NH ₂)]		
20	BisPtmal	=	[(cis-		
			Pt(mal)(NH ₃)] ₂ (H ₂ N(CH ₂) ₄ NH ₂)]		

As can be seen from the data, the complexes of the present invention maintain high activity in the L-1210/DDP line which is resistant to cisplatin. Furthermore comparison of Example 1 with [Pt(dien)Cl]Cl shows that the monomeric complex is totally inactive, in accord with accepted theories, yet the combination of two monodentate units into a bis(platinum) complex as in Example 1 produces highly active cytotoxic agents.

WO 91/03482

PCT/US90/04966

-22-

TABLE III

Toxicity of Complexes of General Formula [PtCl ₂ LL']					
	L	L'	Geometry	ID ₅₀ (μM)	
				L-1210/O	L-1210/DDP
5	py	py	trans	0.61	0.80
	py	py	cis	8.96	12.5
	py	DMSO	trans	6.62	5.67
	Meim	DMSO	trans	6.57	10.33
10	quin	DMSO	trans	0.4	1.57
	iso-quin	DMSO	trans	2.75	1.17
	NH ₃	DMSO	cis	>20	>20
	NH ₃	DMSO	trans	>20	>20
15	NH ₃	NH ₃	cis	0.2	8.0
	NH ₃	NH ₃	trans	>20	>50

20 where py = pyridine; quin = quinoline; isoquin = isoquinoline; Meim = methylimidazole. The biological assays were performed as previously described by method of Hacker et al. Cancer Research, 1985, 45, 4748. Complexes were all dissolved in 10% DMSO.

25 While the invention has been described in terms of various preferred embodiments, one skilled in the art will appreciate that various modifications, substitutions, omissions, and changes may be made without departing from the spirit thereof.

30 Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims.


WO 91/03482

PCT/US90/04966

-23-

WHAT IS CLAIMED IS:

1. A bis(platinum) complex having the structure:



10 wherein X, Y, Z, X', Y' and Z' are the same or different ligands and are an anionic group selected from the group consisting of halide, pseudohalide, substituted pseudohalide, sulphate, phosphate, phosphonate, nitrate, carboxylate, substituted carboxylate, dicarboxylate and substituted

15 dicarboxylate or a neutral group which may be substituted or unsubstituted and is selected from the group consisting of primary or secondary amines, sulfoxide, phosphine, pyridine, acridine, pyrazole, thiazole, imidazole, benzimidazole, benzothiazole,

20 pyrimidine, purine, quinoline and isoquinoline and A is a diamine or polyamine wherein the primary amine N atoms are coordinated to the Pt atom such that platinum is present as Pt²⁺.

- 25 2. The bis(platinum) complex according to claim 1 wherein A has the formula:

NDE — (C(R⁴)₂)_r — (R⁵)_o — (C(R⁶)₂)_p — NDE

wherein r and p are integers from 1 to 4, inclusive, and o is 0 or 1; R⁴ and R⁶ are the same or different

30 and are hydrogen, lower alkyl, aryl, cycloalkyl, cycloalkenyl, aralkyl, halogen, pseudohalogen, hydroxy, alkoxy, aryloxy, carboxylic ester, or carboxylic acid salt; D and E are the same or different and are

35 hydrogen, lower alkyl, aryl, alkaryl, aralkyl, alkenyl, cycloalkyl, cycloalkenyl, halogen, pseudohalogen, hydroxy, alkoxy, aryloxy, sulphonic acid, or sulphonic

WO 91/03482

PCT/US90/04966


-24-

acid salt; and R⁵ is alkyl, aryl, amino, alkylamino, diamino of the formula:



wherein q is an integer of 1 to 4, inclusive,
5 hydroxyalkyl, alkoxy, sulfur or oxygen.

3. A bis(platinum) complex having the structure:



wherein X, Y, Z, X', Y', and Z' are the same or
15 different ligands and are selected from an anionic group selected from the group consisting of halide, pseudohalide, substituted pseudohalide, sulphate, phosphate, phosphonate, nitrate, carboxylate, substituted carboxylate, dicarboxylate, and substituted
20 carboxylate or a neutral group which may be substituted or unsubstituted and is selected from the group consisting of primary or secondary amine, sulfoxide, phosphine, pyridine, acridine, pyrazole, thiazole, imidazole, benzimidazole, benzothiazole, pyrimidine,
25 purine, quinoline or isoquinoline, and wherein n is an integer of 2 to 20 inclusive.

4. The complex of claim 3 wherein Y and Y' are chlorine and X, X', Z, and Z' are all amines.
30

5. The complex of claim 3 wherein X and X' are chlorine and Y, Z, Y' and Z' are amines.

6. The complex of claim 3 wherein Y and X' are chlorine and X, Z, Y' and Z' are amines.
35

WO 91/03482

PCT/US90/04966

-25-

7. The complex of claim 3 wherein Y, Z, X', and Z' are all chlorine and X and Y' are amines.

5 8. The complex of claim 3 wherein Y, Z and Y' are chlorine and X, X' and Z' are amines.

10 9. The complex of claim 3 wherein Y, Z and Z' are chlorine and X, X', and Y' are amines.


10 10. The complex of claim 3 wherein X, Z and Z' are chlorine and X, X', and Y' are amines.

15 11. The complex of claim 3 wherein X, Z and Y' are chlorine and X, X', and Y' are amines.

20 12. The complex of claim 3 wherein either one of X, Y, and Z or, alternatively, X', Y', and Z' are pyridine, quinoline, pyrimidine, purine, sulfoxide, phosphine; one of X, Y, Z and X', Y' and Z' is NH₃; one of X, Y, and Z is chloride, and one of X', Y' and Z' is chloride.

25 13. The complex of claim 3 wherein X, Z, X', and Z' are all chloride, Y is sulfoxide, pyridine, pyrimidine or purine and Y' is an amine.

30 14. The complex of claim 3 wherein either X, Y, Z, X' Y' or Z' is a carboxylate or substituted carboxylate of the formula:



wherein m is an integer from 0 to 5, inclusive; and the R³ groups are the same or different and is hydrogen, substituted or unsubstituted straight or branched

WO 91/03482

PCT/US90/04966

-26-

alkyl, aryl, alkaryl, alkenyl, cycloalkyl, cycloalkenyl, halogen, pseudohalogen, hydroxy, carbonyl, formyl, nitro, amides, amino, alkoxy, aryloxy, sulphonic acid salt, carboxylic acid ester or carboxylic acid salt.

- 15 15. Cytotoxic complexes of trans geometry which contain at least one planar ligand and have the formula $[PtX_2(L)(L')]$ where L is a planar ligand such as pyridine, quinoline, isoquinoline, acridine, pyrazole, thiazole, imidazole, benzimidazole, benzothiazole, and pyridine-like planar aromatic or pseudoaromatic heterocycles and X is an anionic group, with the proviso that if L' is not the same as L then L' represents a primary or second amine or a sulfoxide.


16. A process for producing a bis(platinum) complex, said process comprising:

- (a) reacting a compound of the formula RNH_3^+Cl wherein RNH_3^+ represents $PtX'Y'Z'A$ wherein Z', Y', Z' and A are as defined in claim 1 with a compound of the formula:



wherein X, Y and Z are as defined in claim 1 in the presence of a base;

- and (b) recovering a compound of the formula:



WO 91/03482

PCT/US90/04966

-27-

17. A pharmaceutical composition comprising a tumor-inhibiting effective amount of a complex according to claim 1 in a pharmaceutically acceptable carrier.

5

18. A method of inhibiting tumor growth in a mammal, said method comprising administering a tumor-inhibiting effective amount of a complex according to claim 1.

10

19. A pharmaceutical composition comprising a tumor-inhibiting effective amount of a complex according to claim 3 in a pharmaceutically acceptable carrier.

15

20. A method of inhibiting tumor growth in a mammal, said method comprising administering a tumor-inhibiting effective amount of a complex according to claim 3.

WO 91/03482

PCT/US90/04966

1/1




FIGURE 1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US90/04966

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): C07F 15/00

US CL.: 514/187, 514/188

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System :

Classification Symbols

US

514/187, 514/188, 514/492, 544/225, 546/4,
546/5, 546/6, 548/103, 556/24, 556/136, 556/137

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched *

III. DOCUMENTS CONSIDERED TO BE RELEVANT **

Category *	Citation of Document, * ¹ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
X	US, A, 4,797,393 Published 10 January 1989, (FARRELL ET AL.), columns 2,3 and 4.	1-20
A	US, A, 4,533,502 Published 06 August 1985, (ROCHON ET AL.)	1,17-20
A	US, A, 4,560,782 Published 24 December 1985, (PAPAGEORGIOU ET AL.)	1,17-20
A	US, A, 4,565,884 Published 21 January 1986, (ANDRULIS JR. ET AL.)	1-20
A	US, A, 4,730,069 Published 08 March 1988, (KOLAR ET AL.)	1,17-20

* Special categories of cited documents: ¹³

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international
filing date

"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or
other means

"P" document published prior to the international filing date but
later than the priority date claimed

"-T" later document published after the international filing date
or priority date and not in conflict with the application but
cited to understand the principle or theory underlying the
invention

"-X" document of particular relevance; the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step

"-Y" document of particular relevance; the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art.

"-A" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search *

Date of Mailing of this International Search Report *

30 NOVEMBER 1990

22 JAN 1991

International Searching Authority :

Signature of Authorized ONUSUVER PSCC-RO

ISA/US

INTERNATIONAL DIVISION

Fm ARTHUR C. PRESCOTT *Arthur C. Prescott*